



Is Swedish snus associated with smoking initiation or smoking cessation?

H Furberg, C M Bulik, C Lerman, P Lichtenstein, N L Pedersen and P F Sullivan

Tob. Control 2005;14:422-424
doi:10.1136/tc.2005.012476

Updated information and services can be found at:
<http://tc.bmjournals.com/cgi/content/full/14/6/422>

These include:

References

This article cites 18 articles, 4 of which can be accessed free at:
<http://tc.bmjournals.com/cgi/content/full/14/6/422#BIBL>

Rapid responses

You can respond to this article at:
<http://tc.bmjournals.com/cgi/eletter-submit/14/6/422>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Tobacco use](#) (155 articles)
[Smoking cessation](#) (213 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Tobacco Control* go to:
<http://www.bmjournals.com/subscriptions/>

BRIEF REPORT

Is Swedish snus associated with smoking initiation or smoking cessation?

H Furberg, C M Bulik, C Lerman, P Lichtenstein, N L Pedersen, P F Sullivan

Tobacco Control 2005;14:422–424. doi: 10.1136/tc.2005.012476

Nicotine replacement therapies (NRT) are an effective treatment for tobacco dependence, yet most smokers do not quit or remain abstinent. We investigated whether Swedish snus (snuff) use was associated with smoking cessation among males participating in a large population based twin study in Sweden. Snus use was associated with smoking cessation but not initiation. Given that snus delivers comparable nicotine concentrations but carries lesser cancer risk than cigarettes, snus may be a widely used, non-medical form of NRT. Evaluation of the efficacy of snus for smoking cessation should be evaluated in randomised clinical trials.

Tobacco use is the second major cause of death worldwide. Nicotine replacement therapy (NRT) can double smoking cessation rates, but most smokers receiving treatment do not achieve long term abstinence.¹ Additional NRT is needed.

The prevalence of cigarette smoking is notably low in Sweden while the use of snus is increasing. Swedish snus is a moist smokeless tobacco product that contains lower concentrations of cancer-causing tobacco-specific nitrosamines than found in other smokeless tobacco products and cigarettes.² While snus delivers similar concentrations of nicotine, it carries substantially lower risks of cancer than cigarettes.^{3–8} Male smokers in Sweden appear to be using snus as a form of NRT,⁹ despite a lack of data from randomised clinical trials to support its use as a smoking cessation treatment.¹⁰

We investigated whether lifetime snus use was associated with smoking initiation or cessation to gain insight into its potential role as NRT. It is critical to examine snus' potential dual effects on smoking, given the fear that advocating the use of snus might increase smoking, thereby mitigating its utility as NRT.^{11–13}

PARTICIPANTS, METHODS AND RESULTS

Data from the screening across lifespan twin study (SALT) from the population based Swedish Twin Registry were used.¹⁴ The study protocol was reviewed and approved by the ethical committee of the Karolinska Institute, the Swedish Data Inspection Board, and the Institutional Review Board at the University of North Carolina at Chapel Hill.

All subjects provided written informed consent. Briefly, SALT contains detailed data on tobacco use (type, amount, age at first use, patterns of use, and Fagerstrom test for nicotine dependence (FTND)) from telephone interviews completed by 31 425 twins born in Sweden before 1959. Participants who currently or formerly "smoke(d) at parties", "smoke(d) now and then", or "smoke(d) regularly" were considered "ever smokers". Participants who "only tried cigarettes" or "never smoked" were considered "never smokers". We classified participants as "ever snus user" if they currently or formerly used snus "now and then" or

"regularly"; otherwise they were classified as "never snus user". The questionnaire did not specifically describe what was meant by "regular" or "now and then" tobacco use, rather it was up to the participant to interpret and select the type of tobacco user they considered themselves to be. Since the lifetime prevalence of any snus use was only 2.5% among females, we restricted our analyses to males ($n = 14\,932$).

To investigate whether snus use was associated with smoking initiation, we compared men who used snus *before* they started smoking to men who never used snus in relation to any lifetime smoking (ever versus never "regular" or "now and then" cigarette smokers). To address whether snus use was associated with smoking cessation, we compared men who used snus *after* they began smoking to men who never used snus in relation to smoking status at the time of interview (former versus current "regular" or "now and then" smokers).

Odds ratios (OR) and 95% confidence intervals (CI) were estimated with age adjusted logistic regression models and generalised estimating equations to account for clustering of twin pairs. Stratified analyses were performed to examine whether the associations for smoking cessation remained in subgroups of smokers (heavy versus light smokers; high versus low FTND scores).

Table 1 presents the distributions of tobacco use. Of the 14 932 males that participated in SALT, 14 424 (96.6%) had tobacco use data and ages at initiation. Of these men, 9151 (63.5%) reported smoking during their lifetime. Of the smokers, 7880 (86.1%) reported that they smoke(d) "regularly", 669 (7.3%) smoke(d) "now and then", and 602 (6.6%) "smoke(d) at parties". The prevalence of current smoking status was highest for "now and then" smokers (39.2%) as compared with "regular" smokers (34.0%) and "party" smokers (23.1%). The prevalence of any lifetime snus use in SALT was 28.5% ($n = 4119$), the majority of whom used snus regularly ($n = 3704$, (89.9%)).

"Regular" and "now and then" snus use were inversely associated with smoking initiation (table 2). Only 4.1% of men who ever smoked used snus "regularly" before they started smoking, while 18.5% of non-smokers had used snus "regularly". The odds ratio (OR) for "regular" snus use and ever smoking status was 0.2 (95% confidence interval (CI) 0.2 to 0.3). Despite smaller sample sizes, a similar pattern was observed for men who reported they used snus "now and then". Only 0.5% of men who ever smoked used snus "now and then" before they started smoking, while 1.1% of never smokers reported that they used snus "now and then". "Now and then" snus use was also inversely associated with ever smoking status (OR 0.5, 95% CI 0.3 to 0.7), suggesting that men who used snus "regularly" or "now and then" before they began smoking were less likely to ever smoke.

Abbreviations: FTND, Fagerstrom test for nicotine dependence; NRT, nicotine replacement therapy; SALT, screening across lifespan twin study

Table 1 Distributions of cigarette smoking and snus use among males in the Swedish Twin Registry

	Ever smoker (n=9151, 63.5%)						Never smoker
	Ever regular smoker (n=7880, 86.1%)		Ever now and then smoker (n=669, 7.3%)		Ever party smoker* (n=602, 6.6%)		(n=5273, 36.5%)
	Current regular smoker (n=2683, 34.0%)	Former regular smoker (n=5197, 66.0%)	Current now and then smoker (n=262, 39.2%)	Former now and then smoker (n=407, 60.8%)	Current party smoker (n=139, 23.1%)	Former party smoker: (n=463, 76.9%)	
Regular snus use (n=3704, 25.6%)							
Began using snus <i>before</i> cigarettes	61	147	16	38	.	.	NA
Began using snus <i>after</i> cigarettes	324	1701	57	70	.	.	NA
Only used snus, no cigarettes	NA	NA	NA	NA	NA	NA	976
Began using both at same time	47	230	13	24	.	.	NA
Now and then snus use (n=415, 2.9%)							
Began using snus <i>before</i> cigarettes	11	10	2	6	.	.	NA
Began using snus <i>after</i> cigarettes	100	137	13	20	.	.	NA
Only used snus, no cigarettes	NA	NA	NA	NA	NA	NA	60
Began using both at same time	20	22	3	11	.	.	NA
Never used snus (n=10305, 71.5%)	2120	2950	158	238	139	463	4237

*Ever "party" smokers did not contribute ages at onset of tobacco use, so we could not calculate the age they started smoking in relation to snus use.
NA, not available.

Table 2 Age adjusted odds ratios (OR) (95% confidence intervals (CI)) for "regular" and "now and then" snus use in relation to any lifetime cigarette smoking among males in the Swedish Twin Registry

	Ever smoker*		Never smoker		OR (95%CI)
Regular snus use	262†	4.1%	976	18.5%	0.2 (0.2 to 0.3)
Now and then snus use	29†	0.5%	60	1.1%	0.5 (0.3 to 0.7)
Never used snus	5466	95.4%	4237	80.4%	1.0

*"Ever smoker" includes men who reported smoking "regularly" and "now and then". Men who smoked "at parties" were excluded since they did not contribute ages at onset of tobacco use.

†Men who began using snus *before* cigarettes.

"Regular" snus use was associated with smoking cessation among "regular" and "now and then" smokers (table 3). The proportion of former smokers who used snus "regularly" (34.6%) was higher than the proportion of current smokers who used snus regularly (13.7%). The OR for "regular" snus use and former smoking status was 3.7 (95% CI 3.3 to 4.2), indicating that men who used snus "regularly" were over three times more likely to be former smokers than current smokers. No association was observed between "now and then" snus use and smoking cessation (OR 1.1, 95% CI 0.9 to 1.4). Thus, it appears that only "regular" snus use has an impact on smoking cessation.

DISCUSSION

Consistent with recent studies,^{2,9,15-19} we observed that snus use was associated with smoking cessation, not initiation. Our results support the idea that snus is a type of naturalistic and non-medical NRT that smokers in Sweden may be using to enhance smoking cessation efforts.⁹ We acknowledge the cross sectional nature of our data and assert that this correlation is not necessarily causal. Taken together with the

information presented in a recent debate over the potential of snus as a smoking cessation aide,² we suggest that randomised clinical trials are needed to investigate the utility and risks of snus as NRT under controlled conditions.

We are aware that advocating the use of one addictive tobacco product to diminish the harm from another is a controversial issue, particularly as data supporting the use of snus as NRT could enhance the market of the tobacco industry. Clearly, eliminating all forms of tobacco use would have the most beneficial impacts on world health; however, many smokers are unable to achieve lasting smoking cessation. From a harm reduction perspective, should snus be shown to be as effective as or superior in efficacy to existing NRTs without having adverse health consequences, it may represent a more acceptable means by which to reduce tobacco related health burden.

ACKNOWLEDGEMENTS

This work was supported by CA-085839 to PFS. The funding source had no involvement in study design, collection, analysis, interpretation of the data, and was not involved in the decision to submit this paper for publication.

Table 3 Age adjusted OR (95% CI) for smoking cessation (former versus current smokers) by snus use among males in the Swedish Twin Registry

	Former smoker		Current smoker		OR (95% CI)
Regular snus use	1771†	34.6%	381†	13.7%	3.7 (3.3 to 4.2)
Now and then snus use	157†	3.1%	113†	4.1%	1.1 (0.9 to 1.4)
Never used snus	3188	62.3%	2278	82.1%	1.0

*"Former smokers" and "current smokers" includes men who reported smoking "regularly" and "now and then". Men who smoked "at parties" were excluded since they did not contribute ages at onset of tobacco use.

†Men who began using snus *after* cigarettes.

What this paper adds

This brief report presents findings that Swedish snus (oral snuff) is being used as a naturalistic form of nicotine replacement therapy. We suggest that randomised clinical trials be conducted to assess the efficacy of snus as a smoking cessation aide and to evaluate whether any adverse health consequences result.

CONTRIBUTORS

H Furberg conducted the data analysis and wrote the manuscript. C Bulik provided critical revision for the manuscript for important intellectual content. C Lerman assisted in conception and analytic strategy of research question and contributed to manuscript revisions. P Lichtenstein and N Pedersen assisted in conception and analytic strategy of research question, contributed to manuscript revisions and provided the data from the Swedish Twin Registry. P Sullivan funded the analysis, supervised the data analysis and contributed to manuscript revisions. All authors reviewed and approved the final version of the manuscript.

Authors' affiliations

H Furberg, P F Sullivan*, Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

C M Bulik, Department of Psychiatry, University of North Carolina at Chapel Hill

C Lerman, Department of Psychiatry, Transdisciplinary Tobacco Use Research Center, University of Pennsylvania, Pennsylvania, USA

P Lichtenstein, N L Pedersen†, Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden

*Also Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden; †Also Department of Psychology, University of Southern California, California, USA

Competing interest: None of the authors has a conflict of interest that could inappropriately bias the work. In the late 1970s the Council for Tobacco Research funded collection of data used by N Pedersen in her dissertation. P Sullivan had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Correspondence to: Dr Patrick F Sullivan, Department of Genetics, CB#7264, 4109D Neurosciences Research Building, University of North Carolina, Chapel Hill, NC, 27599-7264, USA; pfsullivan@med.unc.edu

Received 12 May 2005

Accepted 10 August 2005

REFERENCES

- 1 **Silagy C**, Lancaster T, Stead L, *et al*. Nicotine replacement therapy for smoking cessation. *Cochrane Database Syst Rev* 2004;(3):CD000146.
- 2 **Foulds J**, Ramstrom L, Burke M, *et al*. Effect of smokeless tobacco (snus) on smoking and public health in Sweden. *Tobacco Control* 2003;**12**:349–59.
- 3 **Lewin F**, Norell SE, Johansson H, *et al*. Smoking tobacco, oral snuff, and alcohol in the etiology of squamous cell carcinoma of the head and neck: a population-based case-referent study in Sweden. *Cancer* 1998;**82**:1367–75.
- 4 **Schildt EB**, Eriksson M, Hardell L, *et al*. Oral snuff, smoking habits and alcohol consumption in relation to oral cancer in a Swedish case-control study. *Int J Cancer* 1998;**77**:341–6.
- 5 **Ye W**, Ekstrom AM, Hansson LE, *et al*. Tobacco, alcohol and the risk of gastric cancer by sub-site and histologic type. *Int J Cancer* 1999;**83**:223–9.
- 6 **Lagergren J**, Bergstrom R, Lindgren A, *et al*. The role of tobacco, snuff and alcohol use in the aetiology of cancer of the oesophagus and gastric cardia. *Int J Cancer* 2000;**85**:340–6.
- 7 **Accortt NA**, Waterbor JW, Beall C, *et al*. Chronic disease mortality in a cohort of smokeless tobacco users. *Am J Epidemiol* 2002;**156**:730–7.
- 8 **Rodu B**, Jansson C. Smokeless tobacco and oral cancer: a review of the risks and determinants. *Crit Rev Oral Biol Med* 2004;**15**:252–63.
- 9 **Ramstrom L**. Patterns of use: a gate leading to smoking or a way out? *Nicotine Tob Res* 2003;**5**:268.
- 10 **Fagerstrom KO**, Schildt EB. Should the European Union lift the ban on snus? Evidence from the Swedish experience. *Addiction* 2003;**98**:1191–5.
- 11 **Tomar SL**. Smokeless tobacco use is a significant predictor of smoking when appropriately modeled. *Nicotine Tob Res* 2003;**5**:571–3.
- 12 **Stratton K**, Shetty P, Wallace R, *et al*. Clearing the smoke: the science base for tobacco harm reduction—executive summary. *Tobacco Control* 2001;**10**:189–95.
- 13 **Haddock CK**, Weg MV, DeBon M, *et al*. Evidence that smokeless tobacco use is a gateway for smoking initiation in young adult males. *Prev Med* 2001;**32**:262–7.
- 14 **Lichtenstein P**, De Faire U, Floderus B, *et al*. The Swedish Twin Registry: a unique resource for clinical, epidemiological and genetic studies. *J Intern Med* 2002;**252**:184–205.
- 15 **Gilljam H**, Galanti MR. Role of snus (oral moist snuff) in smoking cessation and smoking reduction in Sweden. *Addiction* 2003;**98**:1183–9.
- 16 **Ramstrom L**. Snus: part of the problem or part of the solution? *Addiction* 2003;**98**:1198–9, discussion 1204–7.
- 17 **Rodu B**, Stegmayr B, Nasic S, *et al*. Evolving patterns of tobacco use in northern Sweden. *J Intern Med* 2003;**253**:660–5.
- 18 **Kozlowski LT**, O'Connor RJ, Edwards BQ, *et al*. Most smokeless tobacco use is not a causal gateway to cigarettes: using order of product use to evaluate causation in a national US sample. *Addiction* 2003;**98**:1077–85.
- 19 **O'Connor RJ**, Kozlowski LT, Flaherty BP, *et al*. Most smokeless tobacco use does not cause cigarette smoking: results from the 2000 National Household Survey on Drug Abuse. *Addict Behav* 2005;**30**:325–36.